



# ORTHOPEDIC PARTNERS

EST. 1957

Kenneth Paonessa, M.D.  
Michael Halperin, M.D.  
Gabriel Abella, M.D.  
Mohammad Pasha, M.D.  
Tarik Kardestuncer, M.D.  
Ammar Anbari, M.D.  
Nicole Arcand, M.D.  
Scott Stanat, M.D.  
R. Justin Thoms, M.D.

Steven Wei, M.D.  
Jonathan Puposar, M.D.  
Emily Vafek, M.D.  
Nimit Patel, M.D.  
Peter Wagner, M.D.  
Oksana Zhivotenko, D.O.  
Daniel George, M.D.  
Biren Chokshi, M.D.

Erik Frazier, MHA, CEO

Jonathan R. Puposar, MD

## Osteochondral Allograft Transfer (OATS) Surgery

### Post-Operative Instructions

#### Pain Control and Ice

- A pain medication prescription will be provided prior to discharge.
- You may take the prescribed medication as directed.
- You should expect to experience minimal to moderate knee discomfort for several days and even weeks following the surgery.
- Patients often only need prescription narcotics for a few days following the surgery and then can switch to over-the-counter medications such as Tylenol, Advil, or Ibuprofen.
- Ice bags and elevation should be utilized to decrease swelling and pain. Ice should be applied to the knee up to three times a day for 20 minute periods until swelling subsides

#### Wound Care

- If there is bleeding through the bandage, reinforce it with additional dressings for the first 48 hours.
- After 48 hours, remove the bandage. If the incisions are dry (no active bleeding) then you may leave the knee uncovered without a dressing. Please leave the Steri-strips on across the incisions. If the knee is still bleeding, place new sterile dressings over the incisions and change after an additional 24 hours
- Showering is acceptable after 48 hours. You may get the wounds wet but do NOT scrub the incision sites.
- No swimming or tub bathing until cleared by Dr. Puposar at your first post-operative appointment

#### Rehabilitation/ General Post-Op Care

- You will not be allowed to place weight on the operated leg unless notified by Dr. Puposar. Use crutches for mobilization.
- You will be placed in a Continuous Passive Motion (CPM) machine. You have no motion restrictions. You may begin the settings at -5 to 60\* and advance as tolerated. You should be in the machine for a MINIMUM OF 8 HOURS a day
  - 1) If able to tolerate, you should sleep with this device (slow the speed down to facilitate sleeping)
- **Take one 325 mg (full strength) Aspirin daily for 14 days** (unless otherwise instructed or allergic) to prevent blood clots.
- Your first post-op appointment is 10-14 days following surgery. This is usually scheduled at your pre-operative visit. If not, then please call the office to schedule
- Plan at least 2-3 days away from work or school. Utilize this time to decrease swelling and participate in your home exercise program. You may be able to resume work once the pain and swelling resolves (this varies based on job activity).

#### Important Notes

- Peri-incisional numbness should be expected after any skin incision. Even the small incisions used for a knee arthroscopy.
- Recovery from an arthroscopy can be influenced by many factors. Osteoarthritis of the knee may slow the recovery from a knee arthroscopy and ultimately dictates long term function of the knee.

## Early Post-Operative Exercises

*Start the following exercises as soon as you are able. You can begin these in the recovery room shortly after surgery. You may feel uncomfortable at first, but these exercises will speed your recovery and actually diminish your post-operative pain.*

**Repetitions per set: 20**

**Sets of each exercise: 3**

**Sets per day: 3**


**Quad Sets** - Tighten your thigh muscle. Try to straighten your knee. Hold for 5 to 10 seconds.



**Straight Leg Raises** - Tighten the thigh muscle with your knee fully straightened on the bed, as with the Quad set. Lift your leg several inches. Hold for 5 to 10 seconds. Slowly lower.



**Ankle Pumps** - Move your foot up and down rhythmically by contracting the calf and shin muscles. Perform this exercise periodically for 2 to 3 minutes, 2-3x an hour in the recovery room. Continue this exercise until you are fully recovered and all ankle and lower-leg swelling has subsided.



**Knee Straightening Exercises** - Place a small rolled towel just above your heel so that it is not touching the bed. Tighten your thigh. Try to fully straighten your knee and to touch the back of your knee to the bed. Hold fully straightened for five to 10 seconds.



**Bed-Supported Knee Bends** - Bend your knee as much as possible while sliding your foot on the bed. Hold your knee in a maximally bent position for 5 to 10 seconds and then straighten.

